

Proper Size of the Visible Universe in FRW Metrics with Constant Spacetime Curvature

F. Melia*

*Department of Physics, The Applied Math Program,
and Department of Astronomy, The University of Arizona, AZ 85721, USA*

Abstract

In this paper, we continue to examine the fundamental basis for the Friedmann-Robertson-Walker (FRW) metric and its application to cosmology, specifically addressing the question: What is the *proper* size of the visible universe? There are several ways of answering the question of size, though often with an incomplete understanding of how far light has actually traveled in reaching us today from the most remote sources. The difficulty usually arises from an inconsistent use of the coordinates, or an over-interpretation of the physical meaning of quantities such as the so-called proper distance $R(t) = a(t)r$, written in terms of the (unchanging) co-moving radius r and the universal expansion factor $a(t)$. In this paper, we use the five non-trivial FRW metrics with constant spacetime curvature (i.e., the static FRW metrics, but excluding Minkowski) to prove that in static FRW spacetimes in which expansion began from an initial singularity, the visible universe today has a proper size equal to $R_h(t_0/2)$, i.e., the gravitational horizon at half its current age. The exceptions are de Sitter and Lanczos, whose contents had pre-existing positions away from the origin.

PACS numbers: 04.20.Ex, 95.36.+x, 98.80.-k, 98.80.Jk

*John Woodruff Simpson Fellow; Electronic address: fmelia@email.arizona.edu

I. INTRODUCTION

Recent efforts aimed at providing a better understanding of the fundamental basis for the Friedmann-Robertson-Walker (FRW) metric and its application to cosmology have uncovered several previously unrecognized properties relevant to the interpretation of cosmological data. The standard model of cosmology (Λ CDM) is only marginally consistent with these developing theoretical considerations, reflected in the growing tension between its predictions and what is actually observed, both in the cosmic microwave background (CMB) at high redshift and in the matter distribution and Type Ia supernovae in the nearby Universe.

For example, the use of Birkhoff's theorem and its corollary [1] has shown that the Universe possesses a gravitational horizon (with radius R_h) coincident with the better known Hubble sphere emerging empirically from the observed universal expansion [2]. This new insight has allowed us to consider the impact of strictly adhering to the requirements of both the Cosmological principle and Weyl's postulate [3], which together force R_h to always be equal to ct , the distance light could have traveled during a time t since the big bang (see also [4]). Λ CDM agrees with this constraint only partially, oddly very early in the universal expansion close to the Planck time, and more recently, where the various observations are telling us that $R_h(t_0) \approx ct_0$ today—but not in between. However, a closer examination of the data, particularly the anomalies identified in the fluctuations of the CMB—such as the glaring disparity between the observed angular correlation function and that expected in Λ CDM [5, 6, 7]—is providing accumulating evidence that the $R_h = ct$ condition may indeed be maintained at all redshifts, in contrast to the predictions of the standard model.

This fundamental approach to the study of the cosmological spacetime has also allowed us to examine the nature of cosmological redshift z in FRW metrics with constant spacetime curvature. We recently showed that the interpretation of z as due to the ‘stretching’ of space is coordinate dependent [8]. Namely, we proved that redshift may also be calculated solely from the effects of kinematics and gravitational acceleration, suggesting that its dependence on the expansion factor is simply a manifestation of the high degree of symmetry in FRW, and ought not be viewed as evidence in favor of the idea that space itself is expanding. This distinction between an expanding space and a fixed space through which particles are moving is actually quite critical to the expansion dynamics. For example, the commonly held view that an expanding space can move particles—and even light—faster than c is easily shown

to be false when the cosmological redshift is due solely to the aforementioned kinematic and gravitational effects. The misconception of superluminal expansion arises from the inappropriate usage of the coordinates, but when proper distances are used along with proper times to calculate proper velocities, all of the apparent paradoxes and inconsistencies go away.

An equally important outcome of this study has been a greatly improved understanding of how null geodesics behave in FRW, allowing us to better appreciate which sources are actually observable today. We recently confirmed the importance of R_h in delimiting the size of the observable universe [9] by proving that in all cosmologies with an equation-of-state parameter $w \geq -1$, where the pressure p and density ρ are related by the expression $p = w\rho$, no light rays reaching us today could have ever attained a proper distance $R(t)$ greater than $R_h(t_0)$.

The present paper carries this argument further—continuing to broaden our investigation into the fundamental aspects of the FRW metric—by demonstrating that in static space-times, the most distant sources we see today—particularly the CMB—emitted their light at time $(1/2)t_0$ from a proper distance $R_h(t_0/2)$ away. Applied to the CMB, this result may seem paradoxical because the time t_e at recombination was presumably much earlier than $(1/2)t_0$. Needless to say, this issue has itself caused confusion over the years, with many workers believing that light must have therefore traveled a proper distance $c(t_0 - t_e)$ in reaching us. But it is easy to demonstrate that this was not the case and that the difference is merely due to the time dilation between frames moving at relative speeds close to c . Bikwa et al. [9] demonstrated the basis for this effect by direct numerical calculation of the null geodesics. Here, we use FRW metrics with constant spacetime curvature to prove that, when the expansion begins from a singularity, the proper size of the visible Universe is precisely $R_h(t_0/2)$.

II. THE FRW METRICS WITH CONSTANT SPACETIME CURVATURE

The high degree of symmetry afforded by the FRW metric is a direct consequence of the Cosmological principle and Weyl's postulate, which together require that any distance in the cosmos be expressible as the product of an unchanging comoving radius r and a universal expansion factor $a(t)$ depending only on the cosmic time t (see [10] for a pedagogical

description). The Friedmann-Robertson-Walker (FRW) metric for a spatially homogeneous and isotropic three-dimensional space may be written in the general form,

$$ds^2 = c^2 dt^2 - a^2(t)[dr^2(1 - kr^2)^{-1} + r^2(d\theta^2 + \sin^2\theta d\phi^2)] , \quad (1)$$

where θ and ϕ are the corresponding angular coordinates in the comoving frame. The constant k is $+1$ for a closed universe, 0 for a flat, open universe, or -1 for an open universe.

But this simplification comes at a cost, because the so-called proper distance $R(t) = a(t)r$ in these coordinates is not measured using rulers and clocks at rest with respect to an individual observer; instead, $R(t)$ represents a community distance, compiled from the infinitesimal contributions of myriads of observers lined up between the endpoints, all at the same time t (see, e.g., [11]). Of course, there is nothing intrinsically wrong with the usage of $R(t)$ as a measure of distance—but only so long as one does not over-interpret its physical meaning. For example, a source at R with $dR/dt > c$ is not receding “superluminally,” because although c is measured with rulers and clocks at rest with respect to an individual observer, dR/dt is not (see § III below).

In previous applications (see, e.g., [8]), we had demonstrated that a single observer can assess the speed of expansion relative to c only in terms of his proper distance and proper time, both measured on rulers and clocks at rest with respect to himself [8]. Only then is the speed of light invariant—and always equal to c —and a true upper limit to the speed of any object in the cosmos. It is in this context, therefore, that to meaningfully address the question of how big the visible universe is, the most straightforward way is to first find an alternative set of coordinates dx^μ to rewrite the FRW metric in its static form,

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu , \quad (2)$$

where the metric coefficients $g_{\mu\nu}$ ($\mu, \nu = 0, 1, 2, 3$) are independent of time x^0 , because only then can one claim that the distance and time are being measured at rest with respect to the observer. The form of the metric in Eq. (1) clearly does not satisfy this condition because g_{rr} , $g_{\theta\theta}$ and $g_{\phi\phi}$ are all functions of t through the expansion factor $a(t)$. The FRW metrics that can be transformed in this fashion are those with a constant spacetime curvature [8].

As it turns out, there are exactly six such metrics [12], though one of these—the Minkowski spacetime—is highly trivial; in each of the five remaining cases, a transformation of coordinates permits us to write these solutions in static form (see also [13]). In the following sections, we will consider each of these in turn, the Milne Universe, de Sitter space,

anti-de Sitter space, an open Lanczos-type of Universe, and the Lanczos Universe itself. But we shall also learn that de Sitter and Lanczos [14] are quite different from the rest because they do not begin their expansion from a singularity at time $t = 0$. The size of the visible universe for these two cases is therefore revealingly different from that of all the others.

It is important to stress as we proceed through this exercise that although the spacetime curvature is constant in the cases we consider here, it is generally nonzero. This is a crucial point because the result we obtain is not just an artifact of a cosmology without any spacetime curvature (as in the Milne Universe); it is actually independent of what the spacetime curvature happens to be. In other words, static FRW metrics do not simplify the expansion by eliminating the effects of gravity (or dark energy, for that matter). Gravitational effects are present even when the FRW metric is time-independent, as is well known from the Schwarzschild and Kerr spacetimes.

III. THE “EARLIEST” VISIBLE LIGHT

For simplicity, we will assume radial propagation of light throughout the following discussion. With reference to Eq. (2) written in static form, the proper time and proper distance in this frame are, respectively,

$$d\tau = \sqrt{g_{TT}} dT , \quad (3)$$

and

$$d\mathcal{L} = \sqrt{-ds^2} = \sqrt{g_{RR}} dR , \quad (4)$$

where we write $x^\mu = (cT, R, \theta, \phi)$ for the coordinates that render the FRW metric static. One can trivially see that along any null geodesic (for which $ds = 0$), the proper speed of light in this frame,

$$v_\gamma \equiv \sqrt{\frac{g_{RR}}{g_{TT}}} \frac{dR}{dT} , \quad (5)$$

is always equal to c , irrespective of how curved the spacetime happens to be. Reference to “superluminal” motion is often based on the “coordinate speed,” $dR/dT = c\sqrt{g_{TT}/g_{RR}}$ which, however, is not the speed of light measured on rulers and clocks at rest with respect to the observer.

Let us now ask the question “What was the earliest time T_e at which light we are receiving right now could have been emitted?” As shown in [9], the earliest *cosmic* time t_e at which

this light could have been emitted was 0, because all null geodesics linking us to our past actually started at the origin of the coordinates for $t \rightarrow 0$. But this is not true of the time T_e on the individual observer's clocks, because when viewed in terms of the observer's coordinates, the sources were not at the origin when they emitted the light we see today.

In fact, the proper distance traversed by light in reaching us is equal to the proper distance traveled by the source (starting at the origin) before it emitted that light. Therefore,

$$\int_0^{T_e} v_S \sqrt{g_{TT}} dT = \int_{T_e}^{T_0} c \sqrt{g_{TT}} dT , \quad (6)$$

where v_S is the proper speed of the source and T_0 is the present time. Since the maximum proper speed v_S is c , the condition on T_e reduces to the even simpler form

$$\int_0^{T_e} \sqrt{g_{TT}} dT = \int_{T_e}^{T_0} \sqrt{g_{TT}} dT , \quad (7)$$

which has the obvious solution $T_e = (1/2)T_0$. But $T_0 = t_0$, since t is actually the proper time on the clock at rest with respect to us at our location, and we thus infer that the earliest time light observable to the observer could have been emitted must have been $T_e = (1/2)t_0$. In the next section, we will prove this result for each of the static FRW spacetimes with an expansion from an initial singularity, and then we will discuss what it means to have $T_e \gg t_e$ (i.e., $T_e \gg 0$).

IV. DETAILED PROOF

A. The Milne Universe

The Milne universe [15] has no density ($\rho = 0$) and is characterized by a spatial curvature $k = -1$. It therefore corresponds to a simple solution of Einstein's equations with

$$a(t) = ct , \quad (8)$$

in which the scale factor grows linearly with time at a rate equal to the speed of light c . Since the acceleration $\ddot{a}(t)$ is zero in this cosmology, one might expect such a universe to be flat and a mere re-parametrization of Minkowski space. Indeed, Milne built this type of expansion based solely on special relativity, without any constraints imposed by the more general theory.

To cast the FRW metric for the Milne Universe in its static form, we first introduce the co-moving distance variable χ , defined in terms of r according to

$$r = \sinh \chi , \quad (9)$$

which allows us to write

$$ds^2 = c^2 dt^2 - c^2 t^2 [d\chi^2 + \sinh^2 \chi d\Omega^2] \quad (10)$$

where, for simplicity, we have also introduced the notation $d\Omega^2 \equiv d\theta^2 + \sin^2 \theta d\phi^2$. The transformation that brings Eq. (10) into a static (and manifestly flat) form is

$$\begin{aligned} T &= t \cosh \chi \\ R &= ct \sinh \chi , \end{aligned} \quad (11)$$

for then

$$ds^2 = c^2 dT^2 - dR^2 - R^2 d\Omega^2 . \quad (12)$$

Clearly, a null geodesic in this frame has

$$c dT = -dR \quad (13)$$

(for light approaching us along a radius), and therefore

$$R_e = c(T_0 - T_e) , \quad (14)$$

where the proper distance $\mathcal{L}_e$ to the source in the static coordinate system is here identical to R_e because $g_{TT} = g_{RR} = 1$, and T_e and T_0 are the emission and observation times, respectively, in this frame. To be clear, since the metric coefficients $g_{\mu\nu}$ in Eq. (12) are independent of T , the quantities defined in Eq. (11) are now the coordinates for an individual observer, because time T is being measured on clocks at rest in his frame.

Suppose now that at cosmic time t_e , a source at χ_e emits the light reaching the observer today. The corresponding proper time T_e on the observer's clock at the location of the source will be greater than t_e due to the effects of relativistic time dilation. The most distant sources were moving at proper speeds close to c in this frame when they emitted their light at t_e [8], so the time dilation between T_e and t_e for them approaches infinity. For

this exercise, we may even allow $t_e \rightarrow 0$, in order to examine the time at which the CMB was produced. From Eq. (11), we see that when $t_e \rightarrow 0$,

$$\frac{T_e}{t_e} = \cosh \chi_e \gg 1. \quad (15)$$

In this limit, therefore, $\chi_e \rightarrow \infty$, and so we may write

$$\frac{T_e}{t_e} \rightarrow \frac{1}{2} e^{\chi_e}. \quad (16)$$

But we also know from Eq. (10) that a null geodesic satisfies

$$\int_0^{\chi_e} d\chi' = \int_{t_e}^{t_0} \frac{dt'}{t'}, \quad (17)$$

and so

$$\frac{t_0}{t_e} = e^{\chi_e}. \quad (18)$$

A comparison between Eqs. (16) and (18) immediately tells us that, in the limit $t_e \rightarrow 0$,

$$T_e \rightarrow \frac{1}{2} t_0, \quad (19)$$

a simple and beautiful result that confirms our supposition from the previous section, and one that will be repeated with each subsequent cosmology we consider (except for de Sitter and Lanczos, as we have already anticipated).

Thus, in the Milne Universe, the greatest proper distance (defined in Eq. 14) traveled by light reaching the observer today is only $(1/2)ct_0$, though the sources there look very young due to the effects of time dilation between the source frame and that of the observer. In other words, though the CMB may have been produced at cosmic time $t_e \rightarrow 0$ in the co-moving frame, that event occurred at time $T_e = (1/2)t_0$ in the observer's frame, and though the light signal carries information pertaining to those earliest moments in the Universe's expansion history, according to Eq. 14 it has therefore only traveled a proper distance $(1/2)ct_0$ in reaching the observer.

Later in the discussion section we will describe in greater detail what is actually happening, but the basic principle is rather simple—since in Milne the expansion began from a singularity at time $t = 0$, it took the most distant sources a time $(1/2)t_0$ to reach a proper distance $(R_e = ct_0/2)$ from which the most distant light signal could then have reached the observer by time t_0 . This feature is illustrated schematically in Fig. 1. Note that formally, $R_h = c/H$, and since here $H \equiv \dot{a}/a = 1/t$, we may also write this result in the form $R_e = R_h(t_0/2)$. Observationally, we recognize R_h as the Hubble radius.




FIG. 1: Illustration of the maximum proper distance light could have traveled along (dashed) null geodesics in reaching us as the Universe expands through a metric with constant spacetime curvature. At time t in these spacetimes, we see light only from sources that produced their signal after $(1/2)t$, and were located within a sphere of maximum extent R_e . In all static spacetimes beginning their expansion from an initial singularity, including the Milne universe, $R_e = R_h(t_0/2)$, the gravitational horizon at half the time t_0 elapsed since the big bang.

B. Anti de Sitter Space

A Universe with negative mass density and spatial curvature $k = -1$ is known as anti-de Sitter space, due to its negative spacetime curvature. This metric is given by

$$ds^2 = c^2 dt^2 - (cb)^2 \sin^2(t/b) \left[\frac{dr^2}{1+r^2} + r^2 d\Omega^2 \right], \quad (20)$$

where clearly the expansion factor is now $a(t) = cb \sin(t/b)$. The coordinate transformation

$$R = cbr \sin(t/b), \quad (21)$$

and

$$\tan(T/b) = (1 + r^2)^{1/2} \tan(t/b), \quad (22)$$

produces the static form of the metric,

$$ds^2 = \left[1 + \left(\frac{R}{cb} \right)^2 \right] c^2 dT^2 - \left[1 + \left(\frac{R}{cb} \right)^2 \right]^{-1} dR^2 - R^2 d\Omega^2. \quad (23)$$

Now, along a geodesic connecting the emission point r_e at time t_e with the observer at $r = 0$ and time t_0 ,

$$\int_0^{r_e} \frac{dr}{\sqrt{1+r^2}} = \int_{t_e/b}^{t_0/b} \frac{du}{\sin(u)} , \quad (24)$$

which has the solution [8]

$$r_e = \frac{1}{2} \left(\frac{\tan(t_0/2b)}{\tan(t_e/2b)} - \frac{\tan(t_e/2b)}{\tan(t_0/2b)} \right) . \quad (25)$$

Thus, for a source emitting light very early in the Universe's history, i.e., for $t_e \rightarrow 0$,

$$r_e \rightarrow \frac{1}{2} \frac{\tan(t_0/2b)}{t_e/2b} . \quad (26)$$

In this limit, r_e clearly diverges, so the factor $\sqrt{1+r_e^2}$ in the equation for T_e ,

$$\tan(T_e/b) = (1+r_e^2)^{1/2} \tan(t_e/b) , \quad (27)$$

simply becomes r_e . In addition, $\tan(t_e/b)$ in this limit reduces to t_e/b , and so

$$\tan(T_e/b) \rightarrow \frac{1}{2} \frac{\tan(t_0/2b)}{t_e/2b} \frac{t_e}{b} = \tan(t_0/2b) , \quad (28)$$

which again leads to the beautiful result that

$$T_e \rightarrow \frac{1}{2} t_0 . \quad (29)$$

This is the observer's coordinate time at which the most distant sources visible to him at t_0 produced their light in anti de Sitter space.

C. A Lanczos-type of Universe with $k = -1$

The third FRW metric with constant spacetime curvature is what we refer to as a Lanczos-type universe with $k = -1$ (see also [12, 13]). In co-moving coordinates, this metric may be written

$$ds^2 = c^2 dt^2 - (cb)^2 \sinh^2(t/b) \left[\frac{dr^2}{1+r^2} + r^2 d\Omega^2 \right] , \quad (30)$$

where $a(t) = (cb) \sinh(t/b)$, and b is a constant (though clearly not the Hubble constant $H \equiv \dot{a}/a$). The metric may also be written in static form with the transformation

$$R = cbr \sinh(t/b) , \quad (31)$$

and

$$\tanh(T/b) = (1 + r^2)^{1/2} \tanh(t/b) , \quad (32)$$

which together allow us to write the interval in the form

$$ds^2 = \left[1 - \left(\frac{R}{cb} \right)^2 \right] c^2 dT^2 - \left[1 - \left(\frac{R}{cb} \right)^2 \right]^{-1} dR^2 - R^2 d\Omega^2 , \quad (33)$$

identical (in terms of R and T) to the actual Lanczos metric we shall examine below.

As before, we calculate the co-moving radius r_e to a source at time t_e using the geodesic equation

$$\int_0^{r_e} \frac{dr}{\sqrt{1 + r^2}} = \int_{t_e/b}^{t_0/b} \frac{du}{\sinh(u)} , \quad (34)$$

whose solution is

$$\sinh^{-1}(r_e) = \ln(\tanh[t_0/2b]) - \ln(\tanh[t_e/2b]) . \quad (35)$$

That is,

$$r_e = \frac{1}{2} \left(\frac{\tanh(t_0/2b)}{\tanh(t_e/2b)} - \frac{\tanh(t_e/2b)}{\tanh(t_0/2b)} \right) . \quad (36)$$

The most distant sources visible by the observer at time t_0 emitted their light at time $t_e \rightarrow 0$, so their comoving radius is evidently

$$r_e \rightarrow \frac{1}{2} \frac{\tanh(t_0/2b)}{t_e/2b} . \quad (37)$$

As was true for anti de Sitter space, this radius diverges for the earliest times, and therefore $\sqrt{1 + r_e^2} \rightarrow r_e$. Thus,

$$\tanh(T_e/b) \rightarrow \frac{1}{2} \frac{\tanh(t_0/2b)}{t_e/2b} \frac{t_e}{b} = \tanh(t_0/2b) , \quad (38)$$

so once again

$$T_e \rightarrow \frac{1}{2} t_0 . \quad (39)$$

D. de Sitter Space

We next consider the situation with de Sitter which, unlike Milne, is a cosmology in which objects not only recede from each other, but also accelerate under the influence of gravity. However, unlike anti de Sitter space and the open Lanczos universe we have just considered, de Sitter space does not begin its expansion from a singularity.

The de Sitter cosmology [16] corresponds to a universe devoid of matter and radiation, but filled with a cosmological constant whose principal property is the equation of state $p = -\rho$. The FRW metric in this case may be written

$$ds^2 = c^2 dt^2 - e^{2Ht} [dr^2 + r^2 d\Omega^2] , \quad (40)$$

where $k = 0$ and the expansion factor has the specific form

$$a(t) = e^{Ht} , \quad (41)$$

in terms of the Hubble constant H .

Unlike the Milne model, de Sitter space contains mass-energy (in the form of a cosmological constant). The transformation of coordinates that brings the metric (Eq. 40) into its stationary form is as follows:

$$\begin{aligned} R &= a(t)r \\ T &= t - \frac{1}{2H} \ln \Phi , \end{aligned} \quad (42)$$

where

$$\Phi \equiv 1 - \left(\frac{R}{R_h} \right)^2 , \quad (43)$$

and

$$R_h \equiv \frac{c}{H} \quad (44)$$

is the gravitational (or Hubble) radius. With these, the de Sitter metric becomes

$$ds^2 = c^2 \Phi dT^2 - \Phi^{-1} dR^2 - R^2 d\Omega^2 , \quad (45)$$

and notice that all the metric coefficients are now independent of time T . Of course, the chief difference between this case and that exhibited in Eq. (8), is that the de Sitter spacetime is curved, and therefore both g_{TT} and g_{RR} depend on the spatial coordinates [2]. The form of the metric in Eq. (25) is how de Sitter himself first presented his now famous solution.

To be precise, de Sitter does not actually have an initial singularity because $a(t) \rightarrow 1$ as $t \rightarrow 0$. Rather, the transformation exhibited in Eqs. (42)-(44) demonstrates the dependence of R and T only on the time difference between t and an initial time t_i . The form of these equations corresponds to the choice $t_i = 0$. In talking about the proper size of the visible universe in de Sitter (see § V), we therefore necessarily refer to how far sources and light

have moved during a time $t_0 - t_i$. Again, in the expressions that follow, we will adhere to the convention that $t_i = 0$.

As we did with Milne, let us now consider the time T_e at which the source located at $R_e(t_e)$ emits the light we see today, corresponding to the proper time t_e in its own rest frame. It will be useful for us to borrow a result we derived earlier [8], allowing us to express the redshift in the form

$$1 + z = \frac{1}{1 - R_e(t_e)/R_h} . \quad (46)$$

This follows very easily from the well-known formulation (see, e.g., [11])

$$1 + z = \frac{a(t_0)}{a(t_e)} , \quad (47)$$

or

$$1 + z = \exp [H(t_0 - t_e)] . \quad (48)$$

According to Eqs. (42) and (43),

$$T_e = t_e - \frac{1}{2H} \ln \left[1 - \left(\frac{R_e(t_e)}{R_h} \right)^2 \right] , \quad (49)$$

so that

$$T_e = t_e - \frac{1}{2H} \ln \left[1 - \left(\frac{R_e(t_e)}{R_h} \right) \right] - \frac{1}{2H} \ln \left[1 + \left(\frac{R_e(t_e)}{R_h} \right) \right] . \quad (50)$$

This equation may be manipulated further, yielding

$$T_e = t_e - \frac{1}{2H} \ln e^{-H(t_0 - t_e)} - \frac{1}{2H} \ln \left(\frac{1 + 2z}{1 + z} \right) . \quad (51)$$

And this leads to the result we were seeking,

$$T_e = \frac{1}{2}t_0 + \frac{1}{2}t_e - \frac{1}{2H} \ln \left(\frac{1 + 2z}{1 + z} \right) . \quad (52)$$

To find the time T_e in the observer's frame corresponding to the earliest emission of observable light in the universe, we put $t_e \rightarrow 0$ and $z \rightarrow \infty$, so that

$$T_e \rightarrow \frac{1}{2}t_0 - \frac{1}{2H} \ln 2 . \quad (53)$$

Therefore, when $t_0 \gg 1/H$, we obtain the rather remarkable result that, even in this kind of curved spacetime,

$$T_e \rightarrow \frac{1}{2}t_0 . \quad (54)$$

This is what one would expect on the basis of our discussion in the previous section. As we shall see shortly, however, the proper distance associated with this emission time is not the same as that for the previous three cases.

E. The Lanczos (Closed) Universe

The fifth, and final, FRW metric with constant spacetime curvature (other than Minkowski) is known as the Lanczos Universe, described by the metric

$$ds^2 = c^2 dt^2 - (cb)^2 \cosh^2(t/b) \left[\frac{dr^2}{1-r^2} + r^2 d\Omega^2 \right] , \quad (55)$$

where $k = +1$, and the expansion factor is now $a(t) = (cb) \cosh(t/b)$. This solution represents the gravitational field of a rigidly rotating dust cylinder coupled to a cosmological constant. We use the following transformation to render this metric in static form:

$$R = cbr \cosh(t/b) , \quad (56)$$

and

$$\tanh(T/b) = (1 - r^2)^{-1/2} \tanh(t/b) , \quad (57)$$

which together allow us to write the interval as

$$ds^2 = \left[1 - \left(\frac{R}{cb} \right)^2 \right] c^2 dT^2 - \left[1 - \left(\frac{R}{cb} \right)^2 \right]^{-1} dR^2 - R^2 d\Omega^2 . \quad (58)$$

But it doesn't take much to realize that this universe is quite different from the others. For one thing, it's closed ($k = +1$); the others are all open, with $k = 0$ or $k = -1$. Moreover, as in de Sitter, the expansion factor $a(t)$ does not vanish at $t = 0$. Instead, $a \rightarrow cb$, so this universe does not begin its expansion from a singularity, and because the sources all occupied pre-existing positions at $t = 0$, one should therefore expect that an observer can detect the light they emitted all the way back to $T_e = 0$, which we now demonstrate formally.

As shown in [8], the co-moving distance to a source emitting light at t_e is here given by the expression

$$r_e = 2 \left(e^{t_0/b} - e^{t_e/b} \right) \left(1 + e^{(t_0+t_e)/b} \right) \left(1 + e^{2t_0/b} \right)^{-1} \left(1 + e^{2t_e/b} \right)^{-1} , \quad (59)$$

and it is not difficult to show that as $t_e \rightarrow 0$,

$$r_e \rightarrow \tanh(t_0/b) . \quad (60)$$

Therefore as $t_e \rightarrow 0$,

$$\tanh(T_e/b) \rightarrow \cosh(t_0/b) \tanh(t_e/b) \rightarrow 0 , \quad (61)$$

for which $T_e \rightarrow 0$.

V. PROPER SIZE OF AN FRW UNIVERSE

In all the cases we have considered, the coordinate distance R in the static frame actually coincides with the definition of proper radius $R = a(t)r$ in the co-moving frame. We may therefore determine the “proper” size of the visible universe by merely calculating $R_e(T_e)$ at the time of emission T_e .

The Milne universe is unique among the static FRW metrics because it has zero spacetime curvature. Therefore $g_{RR} = g_{TT} = 1$, and R is exactly equal to the light-travel distance cT , as indicated in Eq. (14). Thus, using Eq. (19), we see that for Milne, $R_e = (1/2)ct_0 = R_h(t_0/2)$.

For anti de Sitter space,

$$R_e = cbr_e \sin(t_e/b) , \quad (62)$$

and introducing the limiting form of r_e from Eq. (26), we find that

$$R_e \rightarrow cb \tan(t_0/2b) , \quad (63)$$

which therefore gives, again, $R_e = R_h(t_0/2)$.

A similar result follows for the open Lanczos universe, since in this case

$$R_e = cbr_e \sinh(t_e/b) , \quad (64)$$

and therefore using the limiting form of r_e from Eq. (37), we find that

$$R_e \rightarrow cb \tanh(t_0/2b) . \quad (65)$$

Since in this case $H(t) = 1/b \tanh(t/b)$, we find that $R_e = R_h(t_0/2)$, as was the case for Milne and anti de Sitter.

These three independent cases all demonstrate the principal result of this paper—that in a static FRW metric expanding from a singularity, the earliest signal we can see today was produced when the Universe was half its current age, t_0 (as measured on clocks at rest with respect to us), from a proper distance equal to the size of the gravitational horizon at that time. As such, this formal derivation is fully consistent with the numerical calculations reported in [9].

Let us now compare this fundamental result with de Sitter and Lanczos, for which $a(0) \neq 0$. It is straightforward to see from Eq. (42) that

$$-t_0 H = \ln \left(1 - \left[\frac{R_e}{R_h} \right]^2 \right) , \quad (66)$$

and therefore $R_e \rightarrow R_h$ for $t_0 H \gg 1$. Even though $T_e = t_0/2$, the proper size of the visible de Sitter universe is nonetheless R_h , which is a constant. The fact that $T_e \neq 0$ is entirely due to our choice of following the expansion from a defined initial time $t_i = 0$. But the structure of this universe is independent of time because its expansion is eternally exponentiated. So the most distant sources we can see are always at the gravitational horizon.

There are strong similarities between de Sitter and the final cosmology we have considered—the Lanczos universe—though the gravitational horizon in this case is not constant. We see from Eq. (56) that

$$R_e = c b r_e \cosh(t_e/b) , \quad (67)$$

so evidently

$$R_e(t_e) \rightarrow c b \tanh(t_0/b) = \frac{c}{H(t_0)} = R_h(t_0) . \quad (68)$$

But even though R_h is here not constant, we find that $R_e \rightarrow c b$ for all $t_0/b \gg 1$, which mirrors the situation with de Sitter. The principal difference between these last two cosmologies and all the others, is that whereas the sources in the previous cases first had to travel a proper distance $R_h(t_0/2)$ to reach the edge of the visible universe, the most distant sources in de Sitter and Lanczos were already situated at $R_e = R_h(\infty)$ from the very beginning, and therefore the observer can see their light emitted from that maximal distance at arbitrarily early times.

VI. DISCUSSION AND CONCLUSIONS

To fully understand and appreciate the results we have presented in this paper, one must acknowledge the critical role played by the choice of coordinates in describing the expansion of the Universe. We had already seen an example of this, based on how the choice of frames impacts our interpretation of the cosmological redshift z [8]. We proved earlier that, although z is conventionally calculated directly from the expansion factor $a(t)$, its origin cannot be attributed to an expansion of space when viewed in terms of the FRW metric written in stationary form. We found that z is actually the cosmological version of a lapse function encountered more typically in the context of the Schwarzschild and Kerr metrics. That is, z is simply due to the combined effects of the kinematic expansion and the gravitational

acceleration—but only in terms of the proper velocity, calculated using the proper distance and proper time for an individual observer.

In this paper, we have expanded our study of the fundamental aspects of the cosmic spacetime by using these alternative sets of coordinates to address another issue that sometimes gives rise to confusion and ambiguity: what is the true size of the visible universe? The question itself is fraught with ambiguity because it goes without saying that to measure a size, one must have a precise definition of distance. General relativity is founded on the basic principle that c is invariant and is measured to have the same value for all observers. But what is often overlooked or forgotten is that in order to make the measurements consistent with this tenet, distances and times must be determined with devices at rest with respect to the observer. Only then can he claim that c is an upper limit to all speeds and that light travels at speed c under all circumstances and at all times.

These notions are particularly important to the question we have addressed in this paper, especially for cosmologies that begin their expansion from a singularity at time $t = 0$. The reason for this is rather straightforward. In these cosmologies, all the worldlines of sources we see today started from the same location—very near the same co-moving point we ourselves are now occupying. Clearly, to suggest that the light they emitted has traveled a distance $c(t_0 - t_e) \rightarrow ct_0$ since the big bang is quite non-sensical. The correct statement is that the most distant sources we see today are precisely those moving at close to proper lightspeed, which reached a proper distance $R_h(t_0/2)$ before emitting the light that is just now reaching us at time t_0 .

It is remarkable—though obvious in retrospect—how elegantly and beautifully this simple result emerges from the properties of the FRW metric itself written in stationary form, when we take the limit $t_e \rightarrow 0$ for the time at which the light from the most distant sources was emitted. One of the principal results of our analysis has been the demonstration that even though $t_e \approx 0$ for these sources, the time measured on our clocks at rest with respect to us was actually $T_e = (1/2)t_0$. And now we understand that this effect is entirely due to the time dilation between us and sources receding at proper speed c when they emitted this light.

These conclusions do come with a caveat, however, because most of these results are based on the use of FRW metrics with a constant spacetime curvature, allowing us to find an alternative set of coordinates to write them in stationary form. Without this option, we

would not yet know how to evaluate distances and times in such a way as to demonstrate without any doubt how far sources could have traveled before emitting the light we see today. One ought to expect the proper size of the visible universe to be measurable against the gravitational horizon even in cases where the spacetime curvature is not constant, but only future work can establish this result conclusively.

Acknowledgements: This research was partially supported by ONR grant N00014-09-C-0032 at the University of Arizona, and by a Miegunyah Fellowship at the University of Melbourne. I am particularly grateful to Amherst College for its support through a John Woodruff Simpson Lectureship.

-
- [1] G. Birkhoff, *Relativity and Modern Physics* (Cambridge: Harvard University Press) (1923).
 - [2] F. Melia, *Mon. Not. R. Astron. Soc.* **382**, 1917 (2007).
 - [3] H. Weyl, *Z. Phys.* **24**, 230 (1923).
 - [4] F. Melia, & A. Shevchuk, *Mon. Not. R. Astron. Soc.* **419**, 2579 (2012).
 - [5] D. N. Spergel *et al.* *Astrop. J. Supp.* **148**, 175 (2003).
 - [6] C. J. Copi, D. Huterer, D. J. Schwarz, & G. D. Starkman, *Mon. Not. R. Astron. Soc.* **399**, 295 (2009).
 - [7] F. Melia, *Phys. Rev. D*, submitted (2012a).
 - [8] F. Melia, *Mon. Not. R. Astron. Soc.* **422**, 1418 (2012b).
 - [9] O. Bikwa, F. Melia, & A. Shevchuk, *Mon. Not. R. Astron. Soc.* **421**, 3356 (2012).
 - [10] F. Melia, *Australian Physics* **49**, 83 (2012c).
 - [11] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (Wiley, New York) (1972).
 - [12] H. P. Robertson & T. W. Noonan, *Relativity and Cosmology* (W. B. Saunders, Philadelphia) (1968).
 - [13] P. S. Florides, *Gen. Rel. Grav.* **12**, 563 (1980).
 - [14] V. K. Lanczos, *Z. Phys.* **21**, 73 (1924).
 - [15] E. A. Milne, *Zeitschrift für Astrophysik* **6**, 1 (1933).
 - [16] W. de Sitter, *Proc. Akad. Wetensch. Amsterdam* **19**, 1217 (1917).